



## Trabajo Original

## Valoración nutricional

### Malnutrition in hospitalized patients: results from La Rioja

### *Desnutrición en pacientes hospitalizados: resultados en La Rioja*

Ángela Martín-Palmero<sup>1,2</sup>, Andra Serrano-Pérez<sup>1</sup>, M.<sup>a</sup> José Chinchetru-Ranedo<sup>2</sup>, Alejandro Cámara-Balda<sup>2</sup>, M.<sup>a</sup> Ángeles Martínez-de-Salinas-Santamaría<sup>2</sup>, Gonzalo Villar-García<sup>2</sup> and M.<sup>a</sup> del Mar Marín-Lizárraga<sup>2</sup>

<sup>1</sup>Unidad de Nutrición Clínica y <sup>2</sup>Servicio de Endocrinología y Nutrición. Hospital San Pedro de Logroño. La Rioja, Spain

## Abstract

**Background:** There is a high malnutrition prevalence in hospitalized patients.

**Aim:** To determine the malnutrition prevalence in hospitalized patients of La Rioja Community (Spain) when evaluated with different screening/evaluation tools and its relationship with hospital stay and mortality.

**Methods:** Cross sectional observational study of hospitalized adult patients (age > 18 years old) from medical and surgical departments that underwent within 72 h of their admission a nutritional screening with Malnutrition Universal Screening Tool (MUST), Nutritional Risk Screening (NRS) 2002, Controlling Nutritional Status (CONUT) y Subjective Global Assessment (SGA).

**Results:** 384 patients (273 medical and 111 surgical) were evaluated. Almost fifty percent of them were considered malnourished independently of the screening/assessment tool used. High concordance was found between SGA and NRS-2002 ( $k = 0.758$ ). Malnourished patients had a longer hospital stay than those well-nourished (9.29 vs. 7.10 days;  $p = 0.002$ ), used a greater number of medicines (9.2 vs. 7.4;  $p = 0.001$ ) and underwent a higher number of diagnostic tests (16.4 vs. 12.5;  $p = 0.002$ ).

**Conclusions:** Half of the hospitalized patients in the medical and surgical department of La Rioja are malnourished. This is associated with a longer hospital stay, higher use of medicines, diagnostics tests and greater mortality. Malnutrition could be detected with easy screening tools to treat it appropriately.

#### Key words:

Hospital malnutrition.  
Nutritional status.  
Nutritional screening.  
Nutritional  
assessment.

## Resumen

**Antecedentes:** existe una alta prevalencia de malnutrición en los pacientes hospitalizados.

**Objetivo:** conocer la prevalencia de la desnutrición en la comunidad de La Rioja con distintos métodos de cribado/valoración nutricional y su relación con la estancia media y mortalidad.

**Métodos:** estudio transversal de 384 pacientes mayores de 18 años (273 pacientes en servicios médicos y 111 en servicios quirúrgicos), a los que se les realizó en las primeras 72 horas del ingreso un cribado/valoración nutricional con MUST (*Malnutrition Universal Screening Tool*), NRS-2002 (*Nutritional Risk Screening*), CONUT (*Controlling Nutritional Status*) y VGS (*Subjective Global Assessment*).

**Resultados:** la desnutrición fue observada en más del 50% de los pacientes independientemente del método de cribado/valoración nutricional utilizado. Existe una fuerte concordancia entre la VGS y el NRS-2002. La desnutrición se relaciona con aumento de la estancia hospitalaria (9,29 vs. 7,10 días;  $p = 0,002$ ), mayor consumo de fármacos (9,2 vs. 7,4;  $p = 0,001$ ) y mayor consumo de pruebas diagnósticas durante la estancia hospitalaria (16,4 vs. 12,5;  $p = 0,002$ ).

**Conclusiones:** uno de cada dos pacientes hospitalizados en la comunidad de La Rioja está desnutrido. La desnutrición se relaciona con aumento de la estancia hospitalaria, aumento de la utilización de pruebas diagnósticas, mayor consumo de medicamentos y aumento de la mortalidad. La desnutrición puede ser detectada con medidas de cribado nutricional sencillas que permitirían tratarla adecuadamente.

#### Palabras clave:

Desnutrición  
hospitalaria. Estado  
nutricional. Cribado  
nutricional. Valoración  
nutricional.

Received: 10/08/2016  
Accepted: 09/01/2017

Martín-Palmero A, Serrano-Pérez A, Chinchetru-Ranedo MJ, Cámara-Balda A, Martínez-de-Salinas-Santamaría MA, Villar-García G, Marín-Lizárraga MM. Malnutrition in hospitalized patients: results from La Rioja. Nutr Hosp 2017;34:402-406

DOI: <http://dx.doi.org/10.20960/nh.458>

#### Correspondence:

Ángela Martín-Palmero. Unidad de Nutrición Clínica.  
Servicio de Endocrinología y Nutrición. Hospital San  
Pedro. C/ Piqueras, nº 98. 26006 Logroño, La Rioja.  
Spain  
e-mail: [anmartinpalmero@gmail.com](mailto:anmartinpalmero@gmail.com)

## INTRODUCTION

Malnutrition is highly prevalent in hospitalized patients. Butterworth et al. in 1974, were the first in talking about malnutrition in these patients (1). In the same year, Bistran et al. reported in two different studies that half of the hospitalized patients (44% medical, 50% surgical) have certain degree of malnutrition (2,3).

Surprisingly, since then the malnutrition percentages reported in last four decades in hospitalized patients has been the roughly the same. Percentages go from 30 to 60%, in studies performed in Spain (4-8) or other parts of the world (9-15).

Malnourished hospitalized patients are at high risk of infections (16), falls (17) and pressure ulcers (18). Malnutrition is also associated with a higher mortality (19,20), higher hospital costs caused by a longer stay higher readmission rates and greater health care services utilization (21).

Consistent data have demonstrated that nutritional treatment with diet modification including oral nutritional supplements (ONS), enteral or parenteral nutrition has consistently demonstrated a beneficial effect on clinical and economical outcomes. These include improvement of the nutritional status (22), muscular mass, strength and performance, morbidity (23) quality of life (24) and mortality. Health care costs decrease is a consequence of a shorter in hospital stay (25), decrease readmission rate (26) and use of health care services in general (27).

Despite this high prevalence, malnutrition in hospitalized patients is a under-diagnosed and undertreated problem (28). Even more, 60% of them worsen their nutritional status during their admission (29).

In the PREDYCES study (30), a multicentric Spanish study, 1 out of 4 inpatients were malnourished, a condition that is associated with a longer hospital stay and with an increased health care costs especially in those that developed malnutrition during their admission.

European authorities have recognized that disease associated malnutrition is a European health problem (Praha Declaration June 11<sup>th</sup> 2009), recommending specific directives to prevent this problem and avoid the morbi-mortality associated to it (31). This directives have been integrated in the European Strategic Health Program UE 2008-2013 (32).

A systematic malnutrition screening on admission and during hospital stay would be desirable aiming at improving clinical and economical outcomes.

The present study was planned to explore the nutritional status of medical and surgical patients admitted to a tertiary university hospital in La Rioja, Spain.

## METHODS

This is a cross sectional study of a prospectively recruited population of a randomly sample of adult inpatients (age > 18 years old) admitted to medical and surgical wards of the Hospital Universitario de San Pedro in Logroño (La Rioja, Spain), from February to June 2011. Sample size was estimated in 384 patients con-

sidering 30-60% malnutrition prevalence with a 95% confidence interval and a precision of 5%.

Nutritional evaluation was performed within the first 72 hours of patient's admission by qualified personnel (certified dietitian). Every day patients were randomly selected using a table of aleatory numbers from the admission list.

Exclusion criteria were patients with an expected hospital stay of less than 72 hours, those with a severe disease with an imminent expected death, patients admitted to Psychiatry, Obstetrics, Ophthalmology, Intensive Care Unit, Pediatrics or those that refused to participate in the study.

Social and demographic data was registered in each patient including diagnosis and number of admissions in the previous year and laboratory lab results. Three different nutritional screening tools were used: Malnutrition Universal Screening Tool (MUST), Nutritional Risk Screening (NRS-2002) and Nutritional Control (CONUT for its spelling in Spanish), using another nutritional evaluation tool: Subjective Global Assessment (SGA). On discharge, number of diagnostic tests performed, days of hospital stay, place of discharge and mortality were registered in all patients.

## STATISTICAL ANALYSIS

Results are presented as mean and standard deviation for quantitative variables and percentages for qualitative variables. Normality was tested in quantitative variables with the Kolmogorov-Smirnov test.

Difference between quantitative variables with normal distribution was evaluated with Student t test or Mann-Whitney U test for those with non-parametric distribution. Difference between quantitative variables was evaluated with  $\chi^2$ . Comparison between different nutritional screening tools was explored with  $\chi^2$  and ANOVA. Concordance between nutritional screening tools was evaluated with kappa index. A statistical significance level of  $p < 0.05$  was determined. The statistical package IBM SPSS Statistics 21 was used for the statistical analysis.

The study was approved by the Clinical Research Ethical Committee of La Rioja and all the participants signed the informed consent.

## RESULTS

Three hundred and eighty four inpatients were evaluated, 71.7% admitted to medical wards and 28.9% to surgical wards. Fifty percent of them were men and their mean age was  $64.6 \pm 16.5$ . Eighteen percent of patients had a previous diagnosis of diabetes mellitus, 42% of high blood pressure, 21% of dyslipemia and 16% had obesity.

Table I shows the cause of admission being neoplastic the most frequent one in 16.7%.

Mean body mass index (BMI) was  $25.7 \text{ kg/m}^2$ , 6.2% of patients had a BMI < 18.5; 33.7% have overweight and 15.7% obesity. Mean percentage of weight loss from regular weight was

**Table I. Diagnosis on admission**

Diagnosis on admission (n = 384)	(%)
Neoplastic	16.7
Acute GI pathology	15.1
Respiratory failure/infection	12.8
Others	12
Scheduled mayor surgery	8.1
Cardiovascular disease	7
Urinary pathology/infection	6
Cerebrovascular disease	4.9
Scheduled major abdominal surgery	3.6
Scheduled other major surgery	3.4
CNS/Cognitive impairment	2.3
Weight loss	2.1
Metabolic decompensation	1.8
Nonscheduled other major surgery	1.8
Nonscheduled major abdominal surgery	1.6
Peripheral vascular disease	0.8


CNS: central nervous system; GI: gastrointestinal.

4.8 ± 5.8. The mean hospital-stay was 8.7 ± 6.8 days and 51% of patients had an admission in the previous 12 months.

Results of the nutritional screening using 3 different nutritional screening tools are shown in figure 1.

All patients underwent a SGA showing that 44% of them were well nourished, 38% mildly malnourish and 18% severely malnourished.

Concordance between SGA and the other screening tools was explored showing a good kappa index with NRS-2002 ( $k = 0.758$ ), moderate with MUST ( $k = 0.422$ ) and weak with CONUT ( $k = 0.340$ ).

**Figure 1.**

Malnutrition prevalence according to the different nutritional screening tools.

Malnutrition according to the cause of admission was 23.4% in neoplastic patients, 18.2% in those with acute GI pathology and 15.4% in those with acute respiratory infection/failure.

No statistical significant gender differences were found in malnutrition. A higher percentage of patients older than 70 years old was malnourish (70.4 vs. 45.7%,  $p < 0.05$ ) and higher in medical than in surgical services (63.1 vs. 38.7%,  $p < 0.05$ ).

Malnourished patients had a mean age 10 years higher than those well-nourished, weighted 11 kg lower and have a BMI 4 points lower. Mean weight loss of malnourished patients were 4.9 kg vs. 0.7 kg of those well-nourished. Statistical significant differences were found also in other anthropometric measurements: tricipital skinfold (TSF), arm circumference (AC) and calf circumference (CC) (Table II).

Malnourished patients also had lower values of albumin, prealbumin, transferrin, cholesterol and CRP (Table III).

Sixty two percent of the malnourished patients had an admission in the previous year compared with only 45.5% of well-nourished ones. Mean hospital stay was 2 days higher in those malnourished (9.29 vs. 7.10,  $p < 0.05$ ). During their hospital stay the number of diagnostic procedures in malnourished patients was higher compared with well-nourished ones ( $16.4 \pm 12.6$  vs.  $12.5 \pm 10.6$ ,  $p < 0.005$ ) and they used a greater number of medicines (9.2 vs. 7.4;  $p = 0.001$ ) (Table IV).

Being malnourished increased the hospital length of stay by 1.8 days (IC ± 95%: 0.3-3.2;  $p < 0.05$ ) in a regression model adjusted by age, gender, weight on admission, BMI, weight loss and diagnosis on admission.

Ninety two percent of the well-nourished patients went home at the time of discharge compared with 77% of those malnourished. A higher percentage of malnourished patients were discharged to a chronic health facility compared with those well-nourished (7 vs. 0.6%,  $p < 0.05$ ). Three percent of the malnourished patients died during their hospital stay compared with only 0.6% of those well-nourished ( $p = 0.06$ ).

Only 17.7% of the malnourished patients received nutritional support during their hospital stay.

**Table II. Anthropometric parameters according to their nutritional state (SGA)**

	Malnourished n = 215	Well-nourished n = 169	
	Mean ± SD	Mean ± SD	p
Weight (kg)	64.61 ± 14.55	75.71 ± 15.55	< 0.001
BMI (kg/m <sup>2</sup> )	23.99 ± 5.11	27.82 ± 5.73	< 0.001
Weight loss (kg)	4.9 ± 5.5	0.7 ± 2.7	< 0.001
TSF (mm)	13.24 ± 6.15	16.43 ± 6.033	< 0.001
AC (cm)	28.49 ± 4.53	31.44 ± 4.04	< 0.001
CC* (cm)	33.11 ± 4.05	36.47 ± 3.60	< 0.001

\*In > 65-years-old. SGA: subjective global assessment; BMI: body mass index; TSF: triceps skinfold; AC: arm circumference; CC: calf circumference.

**Table III.** Laboratory parameters according to their nutritional state (SGA)

	<b>Malnourished n = 215</b>	<b>Well-nourished n = 169</b>	
	<b>Mean <math>\pm</math> SD</b>	<b>Mean <math>\pm</math> SD</b>	<b>p</b>
Albumin (g/dl)	3.5 $\pm$ 0.63	3.97 $\pm$ 0.58	< 0.001
Prealbumin (g/dl)	14.38 $\pm$ 6.78	19.6 $\pm$ 7.28	< 0.01
Transferrin (mg/dl)	186,1 $\pm$ 62	220.8 $\pm$ 56	< 0.01
Cholesterol (mg/dl)	152.79 $\pm$ 42.151	175 $\pm$ 44	< 0.01
Lymphocytes (cell/ml)	1408 $\pm$ 1532	1595 $\pm$ 757	NS
CRP (mg/L)	71.79 $\pm$ 85.80	45.1 $\pm$ 66.1	0.001

SGA: subjective global assessment (SGA); CRP: C-reactive protein.

**Table IV.** Main outcomes

	<b>Malnourished n = 215</b>	<b>Well-nourished n = 169</b>	
	<b>Mean <math>\pm</math> SD</b>	<b>Mean <math>\pm</math> SD</b>	<b>p</b>
Length of stay (day)	9.2 $\pm$ 6.8	7.1 $\pm$ 4.5	< 0.001
Number of diagnostic tests	16.4 $\pm$ 13.5	12.5 $\pm$ 9.4	< 0.001
Medication number	9.2 $\pm$ 4.5	7.4 $\pm$ 3.7	< 0.001

## DISCUSSION

Malnutrition is frequent in hospitalized patients causing a negative impact on mobility and mortality with the expected increase in health care associated costs.

Disease and malnutrition have a two way relationship. While disease could cause malnutrition, malnutrition negatively affects the main outcomes of the disease closing a vicious cycle (33).

La Rioja is an autonomous community with centralized sanitary system with only one tertiary university hospital where the present surveillance was performed.

The study sample of 384 hospitalized patients is representative of the acute hospitalized patients of this autonomous community.

Using the SGA evaluation, malnutrition was found in 56% of the hospitalized patients (38% with moderate and 18% with severe) while using any of the other screening tools (MUST, NRS-2002 y CONUT) malnutrition prevalence was always above 50%. If we compared the malnutrition prevalence found in the present work with the one found in the PREDYCES study (30) including a large Spanish sample of hospitalized patients, the figures found in La Rioja are two times higher indicating that one out of two patients are malnourished in La Rioja while one out of four in the rest of Spain. The present study confirms previous findings from a study performed in 2001 that found only 12.5% of hospitalized patients in La Rioja have a normal nutritional state on discharge, with 55% of them with mild malnutrition and 28.3% severe malnutrition (34). The prevalence of disease-associated malnutrition in other Spanish areas ranges from 24 to 66% (4,5,7). If we compared the present

study findings with the ones from Latin American countries, the figures are quite similar, again with a prevalence of disease associated malnutrition of approximately 50% in the latter (10). The British Association for Parenteral and Enteral Nutrition (BAPEN) estimate that around 34% of the patients admitted to a hospital have some degree of disease associated malnutrition (35). These percentages are similar to those found in other European countries (36,37). The differences found between these studies could be related to study populations and methodology to define malnutrition in each cohort.

The present study also confirmed other studies findings indicating that a higher percentage of malnutrition is found in those patients older than 70 years old (8,30).

We also confirmed that those patients classified as malnourished have a 2.19 days longer hospital stay than those classified as well nourished (4,5,8,27,29) supporting the importance of detecting this problem on admission to avoid important health care associated costs. Then, the detection of malnutrition on admission with easy screening tools could contribute indirectly to save health care associated costs (38).

Our study also confirmed that disease associated malnutrition is also associated with higher medication consumption and performing a higher number of diagnostics tests, findings previously described in other studies performed in Spanish populations (7,29). This confirmed the external validity of our findings and again supporting the critical role of the malnutrition screening tools to detect this important issue.

The percentage of malnourished patients that were discharged home was significantly higher in those well-nourished

while a higher percentage of those malnourished needed to be transferred to a chronic health facility. But most importantly, the mortality rate was also significantly higher in the malnourished patients. These findings are all in line with the findings reported in other Spanish as well as North and Latin American studies (4,5,10,27,28,30).

Finally, the present study also showed that SGA evaluation has a good concordance with NRS-2002 and weaker with the CONUT, confirming previous reports that those screening tools including biochemical parameters are not probably the adequate tools to be used in nutritional screening (39).

There are some limitations in the present study. Firstly, this is a single center study, therefore the present findings could not be translated to other populations. Secondly, we only evaluated three screening tools. We did not included Mini Nutritional Assessment (MNA) that is the recommended screening tools for patients older than 65 years old. Thirdly, the present study only included a limited number of surgical patients, but this is a representative sample of the type of patients seen in our hospital.

The most important findings from the present work are:

- Malnutrition is highly prevalent (almost 50% on admission) in La Rioja.
- Of them only 7% received nutritional support.
- We also confirmed that malnourished patients are associated with worse clinical outcomes, higher health care resource utilization, longer hospital stay and higher mortality.

## REFERENCES

1. Butterworth C. The skeleton in the hospital closet. *Nutrition Today* 1974;9:4-8.
2. Bristian BR, Blackburn GL, Hallowell E, Heddl R. Protein status on general surgical patients. *JAMA* 1974;230:858-60.
3. Bristian BR, Blackburn GL, Vitale J, Cochran D, Naylor J. Prevalence of malnutrition in general medical patients. *JAMA* 1976;235:1567-70.
4. Pérez de la Cruz A, Lobo Tamer G, Orduna Espinosa R, Mellado Pastor C, Aguayo de Hoyos E, Ruiz Lopez MD. Malnutrition in hospitalized patients: prevalence and economic impact. *Med Clin (Barc)* 2004;123:201-6.
5. De Luis D, López Guzmán A. Nutritional status of adult patients admitted to internal medicine departments in public hospitals in Castilla y León, Spain - A multi-centre study. *Eur J Intern Med* 2006;17:556-60.
6. Planas M, Audivert S, Pérez-Portabella C, Burgos R, Puiggrós C, Casanellas JM, et al. Nutritional status among adult patients admitted to an university-affiliated hospital in Spain at the time of genome. *Clin Nutr* 2004;23:1016-24.
7. Martínez Olmos MA, Martínez Vázquez MJ, Martínez-Puga E, el Campo Pérez V. Nutritional status study of inpatients in hospitals of Galicia. *Eur J Clin Nutr* 2005;59:938-46.
8. Sanz París A, García JM, Gómez-Candela C, Burgos R, Martín Á, Matia P; Study VIDA group. Malnutrition prevalence in hospitalized elderly diabetic patients. *Nutr Hosp* 2013;28(3):592-9.
9. Sullivan DH, Sun S, Walls RC. Protein-energy undernutrition among elderly hospitalized patients: a prospective study. *JAMA* 1999;281:2013-9.
10. Correia MI, Campos AC; ELAN Cooperative Study. Prevalence of hospital malnutrition in Latin America: the multicenter ELAN study. *Nutrition* 2003;19:823-5.
11. Norman K, Pichard C, Lochs H, Pirlich M. Prognostic impact of disease-related malnutrition. *Clin Nutr* 2008;27:5-15.
12. Kirkland LL, Kashiwagi DT, Brantley S, Scheurer D, Varkey P. Nutrition in the hospitalized patient. *J Hosp Med* 2013;8:52-8.
13. Charlton KE, Nichols C, Bowden S, Lambert K, Barone L, Mason M, et al. Older rehabilitation patients are at high risk of malnutrition: Evidence from a large Australian database. *J Nutr Health Aging* 2010;14:622-8.
14. Agarwal E, Ferguson M, Banks M, Bauer J, Capra S, Isenring E. Nutritional status and dietary intake of acute care patients: Results from the Nutrition Care Day Survey 2010. *Clin Nutr* 2012;31:41-7.
15. Zhang L, Wang X, Huang Y, Gao Y, Peng N, Zhu W, et al. NutritionDay 2010 audit in Jinling hospital of China. *Asia Pac J Clin Nutr* 2013;22:206-13.
16. Schneider SM, Veyres P, Pivot X, Soummer AM, Jambou P, Filippi J, et al. Malnutrition is an independent factor associated with nosocomial infections. *Br J Nutr* 2004;92:105-11.
17. Bauer JD, Isenring E, Torma J, Horsley P, Martineau J. Nutritional status of patients who have fallen in an acute care setting. *J Hum Nutr Diet* 2007;20:558-64.
18. Fry DE, Pine M, Jones BL, Meimban RJ. Patient characteristics and the occurrence of never events. *Arch Surg* 2010;145:148-51.
19. Lim SL, Ong KC, Chan YH, Loke WC, Ferguson M, Daniels L. Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality. *Clin Nutr* 2012;31:345-50.
20. Correia MI, Waitzberg DL. The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through a multivariate model analysis. *Clin Nutr* 2003;22:235-9.
21. Freijer K, Tan SS, Koopmanschap MA, Meijers JM, Halfens RJ, Nuijten MJ. The economic costs of disease related malnutrition. *Clin Nutr* 2013;32:136-41.
22. Cawood AL, Elia M, Stratton RJ. Systematic review and meta-analysis of the effects of high protein oral nutritional supplements. *Ageing Res Rev* 2012;11:278-96.
23. Milne AC, Potter J, Vivanti A, Avenell A. Protein and energy supplementation in elderly people at risk from malnutrition. *Cochrane Database Syst Rev* 2009:CD003288.
24. Starke J, Schneider H, Altheld B, Stehle P, Meier R. Short-term individual nutritional care as part of routine clinical setting improves outcome and quality of life in malnourished medical patients. *Clin Nutr* 2011;30:194-201.
25. Philipson TJ, Snider JT, Lakdawalla DN, Stryckman B, Goldman DP. Impact of oral nutritional supplementation on hospital outcomes. *Am J Manag Care* 2013;19:121-8.
26. Stratton RJ, Hebuterne X, Elia M. A systematic review and meta-analysis of the impact of oral nutritional supplements on hospital readmissions. *Ageing Res Rev* 2013;12:884-97.
27. Sorensen J, Kondrup J, Prokopowicz J, Schiesser M, Krähenbühl L, Meier R, et al.; EuroOOPS Study Group. EuroOOPS: an international, multicenter study to implement nutritional risk cribado and evaluate clinical outcome. *Clin Nutr* 2008;27:340-9.
28. Corkins MR, Guenter P, DiMaria-Ghalili RA, Jensen GL, Malone A, Miller S, et al.; American Society for Parenteral and Enteral Nutrition. Malnutrition diagnoses in hospitalized patients: United States, 2010. *JPEN* 2014;38(2):186-95.
29. Schenker S. Undernutrition in the UK. *Nutr Bull* 2003;28:87e120.
30. Álvarez-Hernández J, Planas Vila M, León-Sanz M, García de Lorenzo A, Celaya-Pérez S, García-Lorda P, et al.; PREDyCES researchers. Prevalence and costs of malnutrition in hospitalized patients; the PREDyCES Study. *Nutr Hosp* 2012;27:1049-59.
31. Commission of the European Communities. "Together for health: a Strategic Approach for the EU 2008-2013". (Accessed November 25, 2010 at [http://ec.europa.eu/health/strategy/policy/index\\_en.htm](http://ec.europa.eu/health/strategy/policy/index_en.htm)).
32. Jeejeebhoy KN. Nutritional assessment. *Nutrition* 2000;16:585-90.
33. Gonzalez Castela L, Coloma Peral R, Ascorbe Salcedo P, Indo Berges O, Rodríguez Carballo B, Martínez Tutor MJ. Current status of the degree of malnutrition in hospitalized patients of the Community of La Rioja. *Nutr Hosp* 2001;16:7-13.
34. Russell A, Elia M; on behalf of British Association for Parenteral and Enteral Nutrition (BAPEN). Nutrition Cribado Survey in the UK and Republic of Ireland 2010 British Association for Parenteral and Enteral Nutrition (BAPEN). Available at: [www.bapen.org](http://www.bapen.org) [Accessed Jan. 24, 2013].
35. Vanderwee K, Clays E, Bocquaert I, Gobert M, Folens B, Defloor T. Malnutrition and associated factors in elderly hospital patients: A Belgian cross-sectional, multi-centre study. *Clin Nutr* 2010;29:469-76.
36. Meijers JM, Halfens RJ, van Bokhorst-de van der Schueren MA, Dassen T, Schols JM. Malnutrition in Dutch health care: prevalence, prevention, treatment, and quality indicators. *Nutrition* 2009;25(5):512-9.
37. Pirlich M, Schutz T, Norman K, Gastell S, Lubke HJ, Bischoff SC, et al. The German hospital malnutrition study. *Clin Nutr* 2006;25:563-72.
38. Kyle UG, Genton L, Pichard C. Hospital length of stay and nutritional status. *Curr Opin Clin Nutr Metab Care* 2005;8:397-402.
39. Guerra-Sánchez L, Martínez-Rincon C, Fresno Lores M. Cribado nutricional en pacientes con insuficiencia cardíaca: análisis de 5 métodos. *Nutr Hosp* 2015;31:890-9.